

**Central Bureau for Astronomical Telegrams
INTERNATIONAL ASTRONOMICAL UNION**

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.

IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)

CBAT@CFA.HARVARD.EDU (science)

URL <http://www.cfa.harvard.edu/iau/cbat.html> ISSN 0081-0304

Phone 617-495-7440/7244/7444 (for emergency use only)

S/2008 S 1

C. C. Porco, on behalf of the Cassini Imaging Science Team, reports the discovery of S/2008 S 1, an object orbiting within Saturn's G ring in the region between Saturn X/XI (Janus/Epimetheus) and Saturn I (Mimas), in two images of the G ring obtained with Cassini's narrow-angle camera taken on 2008 Aug. 15 and separated by ≈ 30 min. Subsequent observations, and a search of earlier images based on a preliminary orbit fit, have provided additional detections of this object, yielding 21 total detections during 2007 June 15–2009 Feb. 20. A high-precision orbit fit based on a numerical integration of the equations of motion shows that the object is being influenced by a 7:6 mean-motion resonance with Mimas; averaged orbital elements for S/2008 S 1: $a = 167500$ km; $e = 0.0002$, $i = 0^\circ.001$, revolution period = 0.80812 day. The integrated orbit shows that the resonance with Mimas causes a long-period (~ 4 yr) oscillation of ~ 4 km in the semi-major axis of S/2008 S 1, with a corresponding oscillation of a few degrees in its mean longitude. S/2008 S 1 is located within a bright arc in the G ring, so this body likely represents a significant source of the small particles found in this ring. If S/2008 S 1 has the same disk-integrated reflectivity as Saturn XXXIII (Pallene), preliminary estimates of the object's total brightness suggest a physical radius for S/2008 S 1 of ~ 250 m.

COMET C/2007 N3 (LULIN)

D. E. Harker, University of California, San Diego; and C. E. Woodward, University of Minnesota, report on spectrophotometric observations of comet C/2007 N3 using the NASA Infrared Telescope Facility (IRTF) telescope (+ MIRS) on Feb. 22.6 UT. The coma was spatially extended at mid-infrared wavelengths, with a circularly symmetric extent of $\approx 6''$ in diameter (3σ surface brightness). Photometry in a $3''.18$ -diameter circular aperture, centered on the peak in the coma surface brightness coincident with the nucleus position, yields fluxes of within the narrow-band IRTF comet-silicate filters of $[8.7 \mu\text{m}] = 3.58 \pm 0.06$ Jy, $[9.8 \mu\text{m}] = 4.98 \pm 0.09$ Jy, $[10.6 \mu\text{m}] = 5.99 \pm 0.03$ Jy, $[11.6 \mu\text{m}] = 6.82 \pm 0.04$ Jy, and $[12.3 \mu\text{m}] = 7.54 \pm 0.06$ Jy. A Planck-function fit to the photometry yields a color temperature of 267 ± 10 K, which is 14 percent higher than that of a blackbody at an equivalent heliocentric distance (1.38 AU). Both the photometric spectral energy distribution and a low-resolution $10\text{-}\mu\text{m}$ spectrum indicate that C/2007 N3 has no silicate emission feature, implying a deficiency of small, submicron-sized silicate grains in the coma.