# Central Bureau for Astronomical Telegrams INTERNATIONAL ASTRONOMICAL UNION

Mailstop 18, Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A. IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions) CBAT@CFA.HARVARD.EDU (science) URL http://www.cfa.harvard.edu/iau/cbat.html ISSN 0081-0304 Phone 617-495-7440/7244/7444 (for emergency use only)

#### $2009 DD_{45}$

R. P. Binzel, M. Birlan, and F. E. DeMeo, Paris Observatory, report on their 0.8- to 2.5- $\mu$ m spectroscopic measurements of 2009 DD<sub>45</sub> (cf. *MPEC* 2009-D80) on Mar. 2.6 UT using the NASA Infrared Telescope Facility 3-m reflector on Mauna Kea. Absorption bands revealed at 1 and 2  $\mu$ m show the characteristics of the S-type class of minor planets. Using the average albedo value of 0.36 for small near-earth objects in this class (Delbo *et al.* 2003, *Icarus* 166, 116), and based on its H magnitude (25.4), the mean diameter is estimated to be 19  $\pm$  4 m. The Apollo-type object passed only 0.000482 AU from the earth on Mar. 2.57 (cf. *MPEC* 2009-E10).

### COMETS C/2008 V4, C/2008 V5, C/2008 V6 (SOHO)

Further to IAUC 9022, additional near-sun presumed comets have been found on SOHO website images, with their "discovery" observations tabulated below — all Kreutz sungrazers except for C/2008 V5 (non-group). C/2008 V4 was very faint (mag  $\sim$  8–8.5) and condensed. C/2008 V5 was perhaps very slightly diffuse but very small (mag  $\sim$  7.5–8). C/2008 V6 was small and condensed (mag  $\sim$  7.5).

Comet	2008	UT	$\alpha_{2000}$	$\delta_{2000}$	Inst.	$\mathbf{F}$	MPEC
C/2008 V4	Nov.	7.063	$14^{^{\mathrm{h}}}\!46\overset{^{\mathrm{m}}}{.}2$	$-17^{\circ}46^{'}$	C2	RM	2009-C46
C/2008 V5			15  04.1		C2	$_{ m JR}$	2009-C47
C'/2008 V6		11.868	15  06.1	$-18\ 57$	C2	$_{ m JR}$	2009-C47

## COMET C/2007 N3 (LULIN)

Corrigendum. On IAUC 9023, line 12, for 267  $\pm$  10 K, read 276  $\pm$  10 K, Further visual total-magnitude and coma-diameter estimates: Feb. 14.19 UT, 5.5, 10′ (B. H. Granslo, Fjellhamar, Norway, 7×50 binoculars); 20.52, 5.0, 20′ (M. Mattiazzo, Castlemaine, Victoria, Australia, 10×50 binoculars); 24.31, 4.7, 20′ (C. S. Morris, Fillmore, CA, U.S.A., 10×50 binoculars); 27.94, 4.9, 25′ (T. Karhula, Virsbo, Sweden, 8×40 binoculars); Mar. 3.14, 5.3, 25′ (J. J. Gonzalez, Leon, Spain, naked eye).

## NOVA IN THE LARGE MAGELLANIC CLOUD 2009

G. Bianciardi, University of Siena, reports the following V magnitudes for this nova (cf. IAUC 9019) from CCD images obtained remotely using a 0.35-m reflector near Sydney, N.S.W.: Feb. 25.717 UT, 14.50; 26.628, 14.72.